Instruction for stool specimen collection

Test Name	Stool routine OB(Chemical method) Parasite OVA(Direct smear) Amoeba(Direct smear) Neutral Fat Rotavirus Ag			Parasite OVA(Concentrated method)		
Container and Sample Volume	Feces Test Bottle (Volume: peanut size)			(Volume:	mination for Parasite thumb size, f of bottle)	
Note	 Before you do a stool routine or OB (chemical) test, avoid the following for 2 to 3 days before the test: red meat, vitamin C, medicines with iron, Aspirin, peroxidase-containing fruits like banana, grapes, radish. Do not do the test during your menstrual period or if you are having bleeding from hemorrhoids. 			None		
Procedure	 Place the plastic holder (such as bag or paper) in the toilet to collect the stool sample. Avoiding contact with toilet water, urine or toilet paper. Open the lid of bottle, use the plastic stick to get some stool, add stool to the bottle and close the lid tightly. During diarrhea use dropper to collect 2 c.c. of stool to the bottle. 					
Note	Our service and opening E-Da Hospital	Monday to Friday 08: 00~21: 30		urday)~12 : 00	Telephone 886-7-6150011 ext.2803	
	E-Da Cancer Hospital E-Da DaChang Hospital	07:00~16:00 08:00~21:00)~12 : 00)~12 : 00	886-7-6150022 ext.6285 886-7-5599123 ext.7216	